

E.P.A.B. Emergency Physicians Advisory Board	Page:	2-4
Medical Control for MedStar System	Effective Date:	July 1, 2008
	Replaces:	All Previous

Anaphylaxis / Allergic Reaction (Continued)

FRO Advanced / Lead Secondary Paramedic		
1. 12 Lead acquisition and transmission (if available) if myocardial involvement suspected		
Localized Reaction	Localized with SOB	Anaphylaxis / Hypotensive
1. Dexamethasone 10mg IVP/IM flush IV line prior to administration to prevent precipitation.	1. Dexamethasone 10mg IVP/IM flush IV line prior to administration to prevent precipitation.	1. Dexamethasone 10mg IVP/IM flush IV line prior to administration to prevent precipitation.
Conditional Primary Paramedic / Primary Paramedic		
Localized Reaction	Localized with SOB	Anaphylaxis / Hypotensive
1. Reassess and transport	1. Reassess and transport	1. If SBP < 90mmHg, additional 250 ml normal saline IVB , may repeat to a grand total of 1000cc. 2. Epinephrine 1:10,000 infusion 0.01-0.02 mg/min mix 0.1 mg in 10 cc normal saline and administer over 5-10 minutes. 3. Dopamine 5-20 mcg/kg/min infusion for persistent hypotension and to maintain a SBP of 90 mmHg. 4. Reassess and transport
BSP Orders		
1. Further fluid therapy orders		